

INTRINSIC KNOTTING AND LINKING OF ALMOST COMPLETE GRAPHS

JESSE CAMPBELL, THOMAS W. MATTMAN, RYAN OTTMAN, JOEL PYZER, MATT RODRIGUES, AND SAM WILLIAMS

ABSTRACT. We introduce new sufficient conditions for intrinsic knotting and linking. A graph on n vertices with at least $4n - 9$ edges is intrinsically linked. A graph on n vertices with at least $5n - 14$ edges is intrinsically knotted. We also classify graphs that are 0, 1, or 2 edges short of being complete partite graphs with respect to intrinsic linking and intrinsic knotting. In addition, we classify intrinsic knotting of graphs on 8 vertices.

1. INTRODUCTION

We say that a graph is intrinsically knotted (respectively, linked) if every tame embedding of the graph in $\mathbb{R}^3$ contains a non-trivially knotted cycle (respectively, pair of non-trivially linked cycles). Robertson, Seymour, and Thomas [RST] demonstrated that intrinsic linking is determined by the seven Petersen graphs. A graph is intrinsically linked if and only if it contains one of the seven as a minor. Since knotless embedding is preserved under edge contraction [NT], work of Robertson and Seymour [RS] shows that a similar, finite list of graphs exists for the intrinsic knotting property. However, determining this list remains difficult. It is known [CG, F1, KS, MRS] that K_7 and $K_{3,3,1,1}$ along with any graph obtained from these two by triangle-Y exchanges is minor minimal with respect to intrinsic knotting. Recently Foisy [F2, F3] has shown the existence of several additional minor minimal intrinsically knotted graphs.

We will say that a graph is *k -deficient* if it is a complete or complete partite graph with k edges removed. In the current article we classify intrinsic knotting on 1- and 2-deficient graphs as well as graphs on 8 or fewer vertices. As each of the intrinsically knotted graphs in these families contains as a minor K_7 , $K_{3,3,1,1}$, or a graph obtained from these by triangle-Y exchanges, we introduce no new examples of minor minimal intrinsically knotted graphs.

On the other hand, we do introduce a new sufficient condition for intrinsic knotting:

Theorem 1. *A graph on n vertices (where $n \geq 7$) with at least $5n - 14$ edges is intrinsically knotted.*

1991 *Mathematics Subject Classification.* Primary 05C10, Secondary 57M15, 05C35.

Key words and phrases. intrinsic knotting, intrinsic linking, spatial graph, complete partite graphs, topological minor.

This work was completed in part through REU's at CSU, Chico during the summers of 2003 to 2005 supported by NSF REU award #0354174 and the MAA's Strengthening Underrepresented Minority Mathematics Achievement Program, with funding from the NSA, the NSF, and the Moody's Foundation. The CSU, Chico Research Foundation provided additional support.

Moreover, we prove a similar condition for intrinsic linking.

Theorem 2. *A graph on n vertices (where $n \geq 6$) with at least $4n - 9$ edges is intrinsically linked.*

Remark: Theorem 2 was conjectured by Sachs [S] in 1984.

Theorems 1 and 2 follow immediately from the following

Theorem 3. *A graph on n vertices (where $n \geq 7$) with at least $5n - 14$ edges has K_7 as a minor.*

Theorem 4. *A graph on n vertices (where $n \geq 6$) with at least $4n - 9$ edges has K_6 as a minor.*

Indeed, K_7 is intrinsically knotted [CG] and if a graph G has an intrinsically knotted minor, then G is intrinsically knotted [NT]. Similarly, as K_6 is intrinsically linked [CG, S], any graph containing it as a minor is likewise intrinsically linked.

Motivated by Sachs's Conjecture, we found a proof of Theorem 4. Recently, we learned that Mader [M1] had proved both Theorems 3 and 4 in 1968! As our proof of Theorem 4 is quite different from Mader's, we include it as Section 2 of this paper.

In Sections 3, 4, and 5 we investigate 0-, 1-, and 2-deficient graphs respectively. In each case we classify the graphs with respect to intrinsic knotting and linking. (The classification of intrinsic knotting of 0-deficient graphs is due to Blain, Bowlin, Fleming, Foisy, Hendricks, and LaCombe [BFHL].) In Section 6 we classify intrinsically knotted graphs on 8 vertices. (This classification was recently independently presented in [BFHL].)

2. PROOF OF THEOREM 4

In this section we prove:

Theorem 4. *A graph on n vertices (where $n \geq 6$) with at least $4n - 9$ edges has K_6 as a minor.*

We begin with some definitions. A **graph** G is a finite vertex set $V(G)$ and edge family $E(G)$, and we mean G to be simple, connected, and undirected. A graph is labeled K_n , and called the **complete graph** on n vertices, if every vertex in the graph is adjacent to every other vertex. A graph H is a **minor** of a graph G if H results from a finite series of edge contractions, vertex deletions, and edge deletions on G . Two adjacent vertices x and y have a **common neighbor** z in G if $xz, yz \in E(G)$.

Proof: (of Theorem 4) We proceed using induction on n , the number of vertices of our graph G . If $n = 6$, then G has at least $4(6) - 9 = 15$ edges. Since K_6 is the only graph on 6 vertices with 15 edges, $G = K_6$.

We now assume that any graph on $6, 7, \dots, n-1$ vertices (with at least $4|V(G)| - 9$ edges) has a K_6 minor. Let G be a graph on n vertices that has $4n - 9$ edges. (Note that if $|E(G)| > 4n - 9$, then G has a minor on $4n - 9$ edges which can be obtained through edge deletions.)

If G has two adjacent vertices with less than four common neighbors, then contracting their edge results in a graph on $n - 1$ vertices with at least $4(n - 1) - 9 = 4(n - 1) - 9$ edges. By induction, G has a K_6 minor.

Now, suppose $\deg(a) = 6$. Consider the graph H on the neighbors of a : v_1, v_2, v_3, v_4, v_5 , and v_6 . Since every vertex v_i must share at least four common neighbors with a , $\forall v_i \in V(H), \deg_H(v_i) \geq 4$. It follows that H has at least $\frac{6(4)}{2} = 12$ edges. However, we know that if $|E(H)| \geq 3|V(H)| - 5 = 3(6) - 5 = 13$, then H contains a K_5 minor [B, Corollary 1.13], implying that G contains a K_6 minor. So, we may assume that H has exactly 12 edges and each vertex has degree 4. There is only one such graph: a degree 4-regular triangulation on six vertices (see left of Figure 1). Notice that H is K_6 less the edges v_1v_2, v_3v_4 , and v_5v_6 . If we add back any of




FIGURE 1. A vertex of degree 6

Let G' be the subgraph formed from G by removing a and its six edges. Then G' has $n - 1$ vertices and $4(n) - 9 - 6 = 4n - 15$ edges. Let $G' \setminus H$ denote the graph obtained by removing the edges of H . We can assume that there is no path in $G' \setminus H$ from v_1 to v_2 , for otherwise, we can contract along the edges of the path to form a minor in G of the form $a + H'$ with $H' = H \cup v_1 v_2$ having 13 edges. That is, G would have a K_6 minor. Similarly, in $G' \setminus H$ we can assume there is no path

between v_3 and v_4 , nor any path linking v_5 and v_6 . This means that each vertex in $V(G') \setminus V(H)$ is connected to at most three of the vertices in $\{v_1, \dots, v_6\}$ by edges in $G' \setminus H$.

This will allow us to divide G' into eight subgraphs that overlap only in the subgraph H . To illustrate this, we will construct a new graph G'' consisting of the disjoint union of the eight subgraphs. Let's first use H to create eight disjoint triangles $H_1, \dots, H_8$, as at right in Figure 1. Note that this entails doubling each edge of H and making four copies of each vertex. The vertices of G'' will be those of $V(G') \setminus V(H)$ along with the vertices of $H_1, \dots, H_8$. So, we've added three extra copies of each of the six vertices in H and, therefore, G'' will have $n-1+3(6) = n+17$ vertices. The edges of G'' will be those of $G' \setminus H$ along with two copies of each edge of H . Thus, G'' will have $4n - 15 + 12 = 4(n+17) - 71$ edges.

Now, the vertices of $H_1, \dots, H_8$ are connected as eight K_3 's and the vertices in $V(G'') \setminus (V(H_1) \cup \dots \cup V(H_8))$ are connected to one another as they were in $G' \setminus H$. To complete the construction of G'' , we need only describe how the edges between vertices of H and vertices in $V(G') \setminus V(H)$ are placed in G'' . For example, let's look at the edges incident on v_1 . Suppose $w \in V(G') \setminus V(H)$ is a neighbor of v_1 . Then w can be connected in $G' \setminus H$ to at most two other vertices of H . For example, if w is also connected to v_3 and v_5 , then in G'' , we will connect w to the vertex v_1^1 in H_1 , because H_1 is the triangle corresponding to the triangle $v_1 v_3 v_5$ in H . If w is connected to at most one other vertex in H , say v_3 , then in G'' , we have the choice of connecting w to H_1 or H_2 as those are the two triangles containing copies of the vertices v_1 and v_3 . We will adopt the convention of connecting it to the H_i of lower index, in this case H_1 . That is, in G'' there will be an edge between w and the vertex v_1^1 in H_1 . Similarly, if w is connected to no other vertex in H , we have four choices, H_1, H_2, H_3, H_4 , of where to connect w in G'' . Again, we will opt for H_1 , the choice of lowest index. In a similar way, each neighbor w of $v_2, \dots, v_6$ is attached to one of the graphs H_i . This placement is unambiguous if w is connected to three vertices of H , it is made among a choice of two H_i if w is connected to only two vertices, and it is made from a choice of four H_i if w is connected to only one vertex in H . When there is a choice, we will use the H_i of least index i among the given options.

This completes the construction of G'' . Let G_i be the connected component of H_i in G'' . Since G was connected, $G'' \subset \cup_{i=1}^8 G_i$. On the other hand, since we are assuming there is no path in $G' \setminus H$ connecting v_1 and v_2 , v_3 and v_4 , or v_5 and v_6 , the G_i must be disjoint. Now, G'' has $n+17$ vertices and $4(n+17) - 71$ edges. By the pigeon hole principle, this means that one of the subgraphs G_i has n_i vertices and at least $4n_i - \lceil 71/8 \rceil = 4n_i - 8$ edges. This means G_i is not simply a triangle, so $n_i \geq 7$ (recall each vertex has degree at least six). Thus, G_i is a proper subgraph of G that has a K_6 minor.

Finally, suppose $\deg(a) = 7$. Let H be the induced graph on the neighbors of a . Again, each vertex must have degree 4 or more and there are fewer than $3(7) - 5 = 16$ edges. There are several such graphs, but only one has no K_5 minor. This time it is a triangulation on seven vertices with two vertices of degree five and five vertices of degree four, and with 10 triangles (see Figure 2). In this case G'' has $m = n + 22$ vertices and $4m - 89$ edges. There must be a subgraph of G'' that has at least $4n_i - \lfloor \frac{89}{10} \rfloor \geq 4n_i - 8$ edges. So, G has a subgraph that has a K_6 minor, implying G has a K_6 minor. $\square$




FIGURE 2. A vertex of degree 7.

3. COMPLETE AND COMPLETE PARTITE GRAPHS

In this section we present a classification of intrinsic linking of 0-deficient graphs, i.e., complete and complete partite graphs. We also include the classification of intrinsic knotting found in [BFHL] for the reader's convenience. We begin with some lemmas.

If we know that a graph G contains a linked (knotted) minor, then G must also be linked (knotted) [NT]. Conversely, if we can realize G as a minor of an unlinked (unknotted) graph, we deduce that G must also be unlinked (unknotted). A useful lemma in this regard shows how we can combine parts of a k -deficient graph. Recall that $K_{n_1, n_2, \dots, n_p}$ denotes the complete p partite graph with n_i vertices in the i th part. Let $K_{n_1, n_2, \dots, n_p} - k$ denote a graph obtained by removing k edges from $K_{n_1, n_2, \dots, n_p}$.

Lemma 5. $K_{n_1+n_2, n_3, \dots, n_p}$ is a minor of $K_{n_1, n_2, \dots, n_p}$. Similarly, every graph of the form $K_{n_1, n_2, \dots, n_p} - k$ has a minor of the form $K_{n_1+n_2, n_3, \dots, n_p} - k$.

Proof: Combining the n_1 and n_2 parts only involves removing edges between vertices in the n_1 part and the n_2 part. Recall that with complete partite graphs, the ordering of the subscripts is not important, so this lemma implies that we can combine any 2 parts to get a minor of the original graph.

Now, suppose we have a complete partite graph with k edges removed and we combine two parts. The result would be a complete partite graph with one fewer part and at most k edges removed.

Furthermore, if, in $K_{n_1, n_2, \dots, n_k} - k$, there are m edges missing between parts n_1 and n_2 , we can see that $K_{n_1, n_2, \dots, n_k} - k$ has a minor of the form $K_{n_1+n_2, n_3, \dots, n_k} - (k - m)$. $\square$

The other main technique we will use in our classifications (aside from Theorems 1 and 2 and Lemma 5) is based on a lemma for intrinsic linking due to Sachs [S] and an analogous result for intrinsic knotting due to [BFHL, OT]. Let $G + H$ denote the join of graphs G and H , i.e., the graph obtained by taking the

union of G and H and adding an edge between each vertex of G and each vertex of H .

Lemma 6 ([S]). *The graph $G + K_1$ is intrinsically linked if and only if G is non-planar*

Lemma 7 ([BFHL, OT]). *The graph $G + K_2$ is intrinsically knotted if and only if G is non-planar*

Surveying the methods that we will use in the rest of the paper, we see that we will not be introducing any new minor minimal graphs for intrinsically knotting. Indeed, if we use Lemma 7 to argue that a graph $H = G + K_2$ is intrinsically knotted, then, G must contain K_5 or $K_{3,3}$ as a minor. It follows that H contains K_7 or $K_{3,3,1,1}$ as a minor. Similarly, Theorem 1 is based on showing that the graph has a K_7 minor. As our only other technique (demonstrating an intrinsically knotted minor) relies on the graphs derived from K_7 and $K_{3,3,1,1}$ by triangle-Y exchanges, our methods cannot introduce new minor minimal examples. In other words, every intrinsically knotted graph that is 0-, 1-, or 2-deficient or a graph on 8 vertices necessarily has as a minor K_7 , $K_{3,3,1,1}$ or a graph obtained from one of these by triangle-Y exchanges.

Theorem 8. *The complete k -partite graphs are classified with respect to intrinsic linking according to Table 1.*

k	1	2	3	4	5	≥ 6
linked	6	4,4	3,3,1 4,2,2	2,2,2,2 3,2,1,1	2,2,1,1,1 3,1,1,1,1	All
not linked	5	$n,3$	3,2,2 $n,2,1$	2,2,2,1 $n,1,1,1$	2,1,1,1,1	None

TABLE 1. Intrinsic linking of complete k -partite graphs.

Remark: The n in the notation $K_{n,3}$ indicates that the property holds for any number of vertices in the first part, i.e., none of the graphs $K_{1,3}$, $K_{2,3}$, $K_{3,3}$, $\dots$ are intrinsically linked. For each k , the table includes minimal examples of intrinsically linked complete k -partite graphs and maximal graphs which are not intrinsically linked. For example, any complete 2-partite graph which contains $K_{4,4}$ as a minor is intrinsically linked. On the other hand, any complete 2-partite graph which is a minor of a $K_{n,3}$ is not linked. Thus, $K_{l,m}$ is intrinsically linked if and only if $l \geq 4$ and $m \geq 4$.

Proof:

Let us demonstrate that the graphs labeled “linked” in Table 1 are in fact intrinsically linked.

Sachs [S], and, independently, Conway and Gordon [CG] proved that K_6 is linked. Any k -partite graph with $k \geq 6$ contains $K_6 = K_{1,1,1,1,1,1}$ as a minor and is therefore linked. For the remaining k , we appeal to work of Robertson, Seymour, and Thomas [RST] who showed that a graph is intrinsically linked if and only if it contains as a minor one of the seven graphs in the Petersen family. In particular, $K_{4,4}$ with one edge removed and $K_{3,3,1}$ are both in this family. By combining parts,

we see that, for $2 \leq k \leq 5$, one of these two is a minor of each of the “linked” graphs in Table 1.

For each of the “not linked” examples in Table 1 which involve a part with a single vertex, the corresponding graph obtained by removing that vertex is planar. Therefore, by Lemma 7, these graphs are not intrinsically linked.

Since a cycle requires at least three vertices, K_5 has no disjoint pair of cycles and is therefore not linked. The remaining “not linked” graphs in Table 1, $K_{n,3}$ and $K_{3,2,2}$, are, respectively, minors of the unlinked graphs $K_{n,2,1}$ and $K_{2,2,2,1}$. $\square$

For the reader’s convenience, we reproduce the classification of knotted partite graphs due to Blain, Bowlin, Fleming, Foisy, Hendricks, and LaCombe [BFHL] as Table 2 below.

k	1	2	3	4	5	6	≥ 7
knotted	7	5,5	3,3,3 4,3,2 4,4,1	3,2,2,2 4,2,2,1 3,3,1,1	2,2,2,2,1 3,2,2,1,1 3,2,1,1,1	2,2,1,1,1,1 3,1,1,1,1,1	All
not knotted	6	4,4	3,3,2 $n,2,2$ $n,3,1$	2,2,2,2 4,2,1,1 3,2,2,1 $n,2,1,1$	2,2,2,1,1 2,2,1,1,1 $n,1,1,1,1$	2,1,1,1,1,1	None

TABLE 2. Intrinsic knotting of k -partite graphs.

4. 1-DEFICIENT GRAPHS

In this section we classify 1-deficient graphs with respect to intrinsic linking and knotting.

Notation: Often, we will have to talk about a particular vertex or part. We will refer to parts alphabetically with capital letters and vertices of those parts with lower case letters. For example, in $K_{4,3,1}$, we will call the part with 4 vertices part A, the part with 3 vertices part B, and the part with 1 vertex part C. An edge between parts A and C would be labeled (a,c).

4.1. Intrinsic linking.

Theorem 9. *The 1-deficient graphs are classified with respect to intrinsic linking according to Table 3.*

Remark:

Note that most of these graphs have different types of edges. For some graphs, removal of one edge (e.g., (a,b)) will result in a non-linked graph while the removal of a different edge (e.g., (b,c)) will result in a linked graph. However, in many cases, removal of any edge will result in the same categorization. In such cases we simply write “-e”. For example, no matter which edge we remove from $K_{3,2,2}$, we will obtain a graph that is not intrinsically linked.

Proof:

Let us demonstrate that the graphs labeled “linked” in Table 3 are in fact intrinsically linked.

k	1	2	3	4	5	6	≥ 7
linked	7-e	4,4-e	4,3,1-e 3,3,2-e 4,2,2-e	2,2,2,2-e 3,2,1,1-(b,c) 4,2,1,1-e 3,3,1,1-e 3,2,2,1-e	2,2,1,1,1-(b,c) 3,1,1,1,1-(b,c) 4,1,1,1,1-e 3,2,1,1,1-e 2,2,2,1,1-e	2,1,1,1,1,1-e	All
not linked	6-e	$n,3$ -e	3,2,2-e $n,2,1$ -e 3,3,1-e	2,2,2,1-e $n,1,1,1$ -e 3,2,1,1-(a,b) 3,2,1,1-(a,c) 3,2,1,1-(c,d)	2,2,1,1,1-(a,b) 2,2,1,1,1-(c,d) 3,1,1,1,1-(a,b) 2,1,1,1,1-e	1,1,1,1,1,1-e	None

TABLE 3. Intrinsic linking of 1-deficient graphs.

By Theorem 2, K_7 -e, $K_{2,2,2,2}$ -e, $K_{3,2,1,1,1}$ -e, $K_{2,2,2,1,1}$ -e, and $K_{2,1,1,1,1,1}$ -e are intrinsically linked. Any 1-deficient graph with seven or more parts contains K_7 -e as a minor and is therefore intrinsically linked.

Recall that $K_{4,4}$ -e is a Petersen graph. By Lemma 5, it's a minor of $K_{4,3,1}$ -e, $K_{4,2,2}$ -e, $K_{4,2,1,1}$ -e, $K_{3,3,1,1}$ -e, $K_{3,2,2,1}$ -e, and $K_{4,1,1,1,1}$ -e.

$K_{3,3,2}$ -e, and, by Lemma 5, $K_{3,2,1,1}$ -(b,c), $K_{2,2,1,1,1}$ -(b,c), and $K_{3,1,1,1,1}$ -(b,c) all contain the Peterson graph $K_{3,3,1}$ as a minor and are therefore intrinsically linked.

k	1	2	3	4	5
knotted	8-e	5,5-e	3,3,3-e 4,3,2-e 4,4,1-e	3,2,2,2-e 4,2,2,1-e 3,3,2,1-e 4,3,1,1-e	2,2,2,2,1-e 3,2,1,1,1-(b,c) 4,2,1,1,1-e 3,3,1,1,1-e 3,2,2,1,1-e
not knotted	7-e	$n,4$ -e	3,3,2-e $n,2,2$ -e $n,3,1$ -e	3,3,1,1-e 2,2,2,2-e 3,2,2,1-e $n,2,1,1$ -e	3,2,1,1,1-(c,d) 3,2,1,1,1-(a,b) 3,2,1,1,1-(a,c) 2,2,2,1,1-e $n,1,1,1,1$ -e

k	6	7	≥ 8
knotted	2,2,1,1,1,1-(b,c) 3,1,1,1,1,1-(b,c) 3,2,1,1,1,1-e 2,2,2,1,1,1-e 4,1,1,1,1,1-e	2,1,1,1,1,1,1-e	All
not knotted	2,2,1,1,1,1-(a,b) 2,2,1,1,1,1-(c,d) 3,1,1,1,1,1-(a,b) 2,1,1,1,1,1-e	1,1,1,1,1,1,1-e	None

TABLE 4. Intrinsic knotting of 1-deficient graphs.

Next, we'll argue that the "not linked" graphs are in fact not intrinsically linked.

k	1	2	3	4
linked	7-2e	5,4-2e	4,3,1- $\{(b, c), e\}$	3,2,1,1- $\{(b_1, c), (b_1, d)\}$
			4,3,1- $\{(a_1, b_1), (a_1, b_2)\}$	3,2,1,1- $\{(b_1, c), (b_2, c)\}$
			4,3,1- $\{(a_1, b_1), (a_1, c)\}$	4,2,1,1- $\{(b, c), e\}$
			3,3,2- $\{(a_1, b_1), (a_1, b_2)\}$	4,2,1,1- $\{(c, d), e\}$
			3,3,2- $\{(a_1, c_1), (a_2, c_1)\}$	4,2,1,1- $\{(a_1, b_1), (a_1, b_2)\}$
			3,3,2- $\{(a_1, c_1), (b_1, c_1)\}$	4,2,1,1- $\{(a_1, b_1), (a_1, c)\}$
			3,3,2- $\{(a_1, b_1), (b_2, c_1)\}$	4,2,1,1- $\{(a_1, c), (a_1, d)\}$
			3,3,2- $\{(a_1, c_1), (b_1, c_2)\}$	3,3,1,1- $\{(b, c), e\}$
			3,3,2- $\{(a_1, c_1), (a_1, c_2)\}$	3,3,1,1- $\{(c, d), e\}$
			4,2,2- $\{(b, c), e\}$	3,3,1,1- $\{(a_1, b_1), (a_1, b_2)\}$
			4,2,2- $\{(a_1, b_1), (a_1, b_2)\}$	3,2,2,1- $\{(b, c), e\}$
			5,3,1-2e	3,2,2,1- $\{(c, d), e\}$
			4,4,1-2e	3,2,2,1- $\{(a, d), e\}$
			4,3,2-2e	3,2,2,1- $\{(a_1, b_1), (a_1, b_2)\}$
not linked	6-2e	4,4-2e n,3-2e	4,3,1- $\{(a_1, b_1), (a_2, b_1)\}$	3,2,1,1- $\{(a, b), e\}$
			4,3,1- $\{(a_1, b_1), (a_2, b_2)\}$	3,2,1,1- $\{(a, c), e\}$
			4,3,1- $\{(a_1, c), (a_2, c)\}$	3,2,1,1- $\{(c, d), e\}$
			4,3,1- $\{(a_1, b_1), (a_2, c)\}$	3,2,1,1- $\{(b_1, c), (b_2, d)\}$
			3,3,2- $\{(a_1, b_1), (a_2, b_2)\}$	4,2,1,1- $\{(a_1, b_1), (a_2, b_2)\}$
			3,3,2- $\{(a_1, b_1), (b_1, c_1)\}$	4,2,1,1- $\{(a_1, c), (a_2, c)\}$
			3,3,2- $\{(b_1, c_1), (b_2, c_2)\}$	4,2,1,1- $\{(a_1, b_1), (a_2, b_1)\}$
			4,2,2- $\{(a_1, b_1), (a_2, b_2)\}$	4,2,1,1- $\{(a_1, b_1), (a_2, c)\}$
			4,2,2- $\{(a_1, b_1), (a_2, b_1)\}$	4,2,1,1- $\{(a_1, c), (a_2, d)\}$
			4,2,2- $\{(a_1, b_1), (a_2, c_1)\}$	3,3,1,1- $\{(a_1, b_1), (a_2, b_2)\}$
			4,2,2- $\{(a_1, b_1), (a_1, c_1)\}$	3,2,2,1- $\{(a_1, b_1), (a_2, b_2)\}$
			3,2,2-2e	3,2,2,1- $\{(a_1, b_1), (a_1, c_1)\}$
			n,2,1-2e	2,2,2,1-2e
			3,3,1-2e	n,1,1,1-2e

TABLE 5. Intrinsic linking of 2-deficient graphs

K_6 (or $K_{1,1,1,1,1,1}$) and $K_{3,3,1}$ are Petersen graphs and therefore minor minimal with respect to intrinsic linking [RST], so K_6 -e and $K_{3,3,1}$ -e are not intrinsically linked by definition of minor minimal.

$K_{n,3}$ -e, $K_{3,2,2}$ -e, $K_{n,2,1}$ -e, $K_{2,2,2,1}$ -e, $K_{n,1,1,1}$ -e, and $K_{2,1,1,1,1}$ -e are all not intrinsically linked before the edge is removed; so, naturally, we can remove an edge and still have a non-intrinsically linked graph.

$K_{3,2,1}$ -(c,d) is equivalent to $K_{3,2,2}$ which is not intrinsically linked.

k	5	6	≥ 7
linked	$2,2,1,1,1-\{(b_1, c), (b_2, c)\}$ $2,2,1,1,1-\{(a_1, d), (b_1, c)\}$ $2,2,1,1,1-\{(b_1, c), (b_1, d)\}$ $3,1,1,1,1-\{(b, c), (c, d)\}$ $4,1,1,1,1-\{(b, c), e\}$ $4,1,1,1,1-\{(a_1, b), (a_1, c)\}$ $3,2,1,1,1-\{(a, c), e\}$ $3,2,1,1,1-\{(b, c), e\}$ $3,2,1,1,1-\{(c, d), e\}$ $3,2,1,1,1-\{(a_1, b_1), (a_1, b_2)\}$ $3,2,1,1,1-\{(a_1, b_1), (a_2, b_1)\}$ $2,2,2,1,1-2e$ $5,1,1,1,1-2e$ $4,2,1,1,1-2e$ $3,3,1,1,1-2e$	$2,1,1,1,1,1-\{(a_1, b), (a_1, c)\}$ $2,1,1,1,1,1-\{(a_1, b), (a_2, b)\}$ $2,1,1,1,1,1-\{(a_1, b), (c, d)\}$ $2,1,1,1,1,1-\{(b, c), (c, d)\}$ $3,1,1,1,1,1-2e$ $2,2,1,1,1,1-2e$	All
not linked	$2,2,1,1,1-\{(a, b), e\}$ $2,2,1,1,1-\{(c, d), e\}$ $2,2,1,1,1-\{(b_1, c), (b_2, d)\}$ $2,2,1,1,1-\{(a_1, c), (b_1, c)\}$ $3,1,1,1,1-\{(a, b), e\}$ $3,1,1,1,1-\{(b, c), (d, e)\}$ $4,1,1,1,1-\{(a_1, b), (a_2, c)\}$ $4,1,1,1,1-\{(a_1, b), (a_2, b)\}$ $3,2,1,1,1-\{(a_1, b_1), (a_2, b_2)\}$ $2,1,1,1,1-2e$	$2,1,1,1,1,1-\{(a_1, b), (a_2, c)\}$ $2,1,1,1,1,1-\{(a_1, b), (b, c)\}$ $2,1,1,1,1,1-\{(b, c), (d, e)\}$ $1,1,1,1,1,1-2e$	None

TABLE 6. Intrinsic linking of 2-deficient graphs (cont.)

The remaining graphs all have a vertex that is connected to every other vertex, and the removal of that vertex results in a planar graph. So, by Lemma 6, none are intrinsically linked. $\square$

4.2. Intrinsic knotting.

Theorem 10. *The 1-deficient graphs are classified with respect to intrinsic knotting according to Table 4.*

Proof:

Let us demonstrate that the graphs labeled “knotted” in Table 4 are in fact intrinsically knotted.

By Theorem 1, K_8-e , $K_{2,2,2,2,1-e}$, $K_{3,2,1,1,1,1-e}$, $K_{2,2,2,1,1,1-e}$, and $K_{2,1,1,1,1,1,1-e}$ are intrinsically knotted. Any 1-deficient graph of eight or more parts will contain K_8-e as a minor and therefore be intrinsically knotted.

$K_{5,5-e}$ is intrinsically knotted as proved in [Sh].

$K_{3,3,3-e}$ and $K_{4,4,1-e}$ have the graph H_9 of [KS] (derived from K_7 by two triangle-Y moves) as a minor and are therefore intrinsically knotted. By Lemma 5, $K_{3,3,2,1-e}$, $K_{3,3,1,1,1-e}$, and $K_{3,2,2,1,1-e}$ all have $K_{3,3,3-e}$ as a minor, so they too are intrinsically knotted.

k	1	2	3
knotted	8-2e	5,5-2e	$3,3,3-\{(a_1, b_1), (a_1, b_2)\}$
			$3,3,3-\{(a_1, b_1), (b_2, c_1)\}$
			$4,4,1-\{(a_1, c), (b_1, c)\}$
			$4,4,1-\{(a_1, b_1), (b_1, c)\}$
			$4,3,2-\{(a_1, b_1), (a_1, b_2)\}$
			$4,3,2-\{(a_1, c_1), (b_1, c_2)\}$
			$4,3,2-\{(a_1, c_1), (b_1, c_1)\}$
			$4,3,2-\{(b_1, c_1), (b_2, c_1)\}$
			$4,3,2-\{(a_1, c_1), (a_1, c_2)\}$
			$4,3,2-\{(a_1, b_1), (b_2, c_1)\}$
not knotted	7-2e	n,4-2e	$4,3,2-\{(b_1, c_1), (b_1, c_2)\}$
			5,4,1-2e
			5,3,2-2e
			4,3,3-2e
			4,4,2-2e
			$3,3,3-\{(a_1, b_1), (b_1, c_1)\}$
			$3,3,3-\{(a_1, b_1), (a_2, b_2)\}$
			$4,4,1-\{(a_1, b_1), (a_1, b_2)\}$
			$4,4,1-\{(a_1, b_1), (a_2, b_2)\}$
			$4,4,1-\{(a_1, c), (a_2, c)\}$
			$4,4,1-\{(a_1, b_1), (a_2, c)\}$
			$4,3,2-\{(a_1, b_1), (a_2, b_1)\}$
			$4,3,2-\{(a_1, b_1), (a_2, b_2)\}$
			$4,3,2-\{(a_1, c_1), (a_2, c_1)\}$
			$4,3,2-\{(a_1, b_1), (b_1, c_1)\}$
			$4,3,2-\{(a_1, b_1), (a_2, c_1)\}$
			$4,3,2-\{(b_1, c_1), (b_2, c_2)\}$
			$4,3,2-\{(a_1, c_1), (a_2, c_2)\}$
			$4,3,2-\{(a_1, b_1), (a_1, c_1)\}$
			3,3,2-2e
			n,2,2-2e
			n,3,1-2e

TABLE 7. Intrinsic knotting of 2-deficient graphs

$K_{4,3,2-e}$ is intrinsically knotted. Depending on which edge is removed, the graph will either have H_9 of [KS] or $K_{3,3,1,1}$ as a minor. Consequently, by Lemma 5, $K_{3,2,2,2-e}$, $K_{4,2,2,1-e}$, $K_{4,3,1,1-e}$, $K_{4,2,1,1,1-e}$, and $K_{4,1,1,1,1,1-e}$ are intrinsically knotted.

Finally, $K_{3,2,1,1,1-(b,c)}$, $K_{2,2,1,1,1,1-(b,c)}$, and $K_{3,1,1,1,1,1-(b,c)}$, all have $K_{3,3,1,1}$ as a minor and are therefore intrinsically knotted.

Now we'll show that the "not knotted" graphs are not in fact intrinsically knotted.

K_7 was shown to be minor minimally intrinsically knotted in [CG] and $K_{3,3,1,1}$ was shown to be minor minimally intrinsically knotted in [F1], so K_7-e (or $K_{1,1,1,1,1,1,1-e}$) and $K_{3,3,1,1-e}$ are not intrinsically knotted.

k	4		5	
knotted	3,2,2,2- $\{(b,c),e\}$	3,3,2,2-2e	3,2,1,1,1- $\{(b_1,c),(b_1,d)\}$	
	3,2,2,2- $\{(a_1,b_1),(a_1,b_2)\}$	5,2,2,1-2e	3,2,1,1,1- $\{(b_1,c),(b_2,c)\}$	
	3,2,2,2- $\{(a_1,b_1),(a_2,b_1)\}$	4,3,2,1-2e	4,2,1,1,1- $\{(c,d),e\}$	
	3,2,2,2- $\{(a_1,b_1),(a_2,c_1)\}$	3,3,3,1-2e	4,2,1,1,1- $\{(b,c),e\}$	
	4,2,2,1- $\{(b,c),e\}$	5,3,1,1-2e	4,2,1,1,1- $\{(a_1,b_1),(a_1,c)\}$	
	4,2,2,1- $\{(c,d),e\}$	4,4,1,1-2e	4,2,1,1,1- $\{(a_1,b_1),(a_1,b_2)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_1,b_2)\}$		4,2,1,1,1- $\{(a_1,c),(a_1,d)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_1,d)\}$		3,3,1,1,1- $\{(b,c),e\}$	
	3,3,2,1- $\{(c,d),e\}$		3,3,1,1,1- $\{(c,d),e\}$	
	3,3,2,1- $\{(a,d),e\}$		3,3,1,1,1- $\{(a_1,b_1),(a_2,b_1)\}$	
	3,3,2,1- $\{(a_1,b_1),(a_1,b_2)\}$		3,2,2,1,1- $\{(a,d),e\}$	
	3,3,2,1- $\{(b_1,c_1),(b_1,c_2)\}$		3,2,2,1,1- $\{(c,d),e\}$	
	3,3,2,1- $\{(a_1,c_1),(b_1,c_1)\}$		3,2,2,1,1- $\{(b,c),e\}$	
	3,3,2,1- $\{(a_1,b_1),(a_2,c_1)\}$		3,2,2,1,1- $\{(d,e),e\}$	
not knotted	3,3,2,1- $\{(a_1,c_1),(a_2,c_1)\}$	3,3,1,1-2e	3,2,1,1,1- $\{(a,b),e\}$	
	3,3,2,1- $\{(a_1,b_1),(a_2,b_2)\}$	2,2,2,2-2e	3,2,1,1,1- $\{(a,c),e\}$	
	4,2,2,1- $\{(a_1,b_1),(a_2,b_1)\}$	3,2,2,1-2e	3,2,1,1,1- $\{(c,d),e\}$	
	4,2,2,1- $\{(a_1,d),(a_2,d)\}$	n,2,1,1-2e	3,2,1,1,1- $\{(b_1,c),(b_2,d)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_2,b_2)\}$		4,2,1,1,1- $\{(a_1,b_1),(a_2,b_2)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_2,c_1)\}$		4,2,1,1,1- $\{(a_1,b_1),(a_2,c)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_2,d)\}$		4,2,1,1,1- $\{(a_1,b_1),(a_2,b_1)\}$	
	4,2,2,1- $\{(a_1,b_1),(a_1,c_1)\}$		4,2,1,1,1- $\{(a_1,c),(a_2,c)\}$	
	3,3,2,1- $\{(a_1,b_1),(b_1,c_1)\}$		4,2,1,1,1- $\{(a_1,c),(a_2,d)\}$	
	3,3,2,1- $\{(a_1,b_1),(a_2,b_2)\}$		3,3,1,1,1- $\{(a_1,b_1),(a_2,b_2)\}$	
	3,3,2,1- $\{(b_1,c_1),(b_2,c_2)\}$		3,2,2,1,1- $\{(a_1,b_1),(a_1,c_1)\}$	
	4,3,1,1- $\{(a_1,b_1),(a_2,b_1)\}$		3,2,2,1,1- $\{(a_1,b_1),(a_2,b_2)\}$	
	4,3,1,1- $\{(a_1,b_1),(a_2,b_2)\}$		2,2,2,1,1-2e	
	4,3,1,1- $\{(a_1,b_1),(a_2,c)\}$		n,1,1,1,1-2e	
4,3,1,1- $\{(a_1,c),(a_2,d)\}$				
4,3,1,1- $\{(a_1,c),(a_2,c)\}$				

TABLE 8. Intrinsic knotting of 2-deficient graphs (cont.)

In [BFHL], the authors show that $K_{4,4}$ is not intrinsically knotted. In fact, the proof really shows that $K_{n,4}$ is not intrinsically knotted, so, naturally, $K_{n,4}-e$ is not intrinsically knotted either. Similarly, $K_{3,3,2}-e$, $K_{n,2,2}-e$, $K_{n,3,1}-e$, $K_{2,2,2,2}-e$, $K_{3,2,2,1}-e$, $K_{n,2,1,1}-e$, $K_{2,2,2,1,1}-e$, $K_{n,1,1,1,1}-e$, and $K_{2,1,1,1,1,1}-e$ are not intrinsically knotted.

$K_{3,2,1,1,1}-\{c,d\}$ is equivalent to $K_{3,2,2,1}$.

The remaining 5 graphs each have 2 vertices connected to every other vertex. If those 2 vertices are deleted, the resulting graph is planar, so by Lemma 7, they are not intrinsically knotted. $\square$

k	6	7	≥ 8
knotted	$2,2,1,1,1,1-\{(b_1,c),(b_2,c)\}$	$2,1,1,1,1,1,1-\{(a_1,b),(a_1,c)\}$	all
	$2,2,1,1,1,1-\{(b_1,c),(b_1,d)\}$	$2,1,1,1,1,1,1-\{(a_1,b),(a_2,b)\}$	
	$2,2,1,1,1,1-\{(a_1,d),(b_1,c)\}$	$2,1,1,1,1,1,1-\{(a_1,b),(c,d)\}$	
	$3,1,1,1,1,1-\{(b,c),(c,d)\}$	$2,1,1,1,1,1,1-\{(b,c),(c,d)\}$	
	$4,1,1,1,1,1-\{(b,c),e\}$	$3,1,1,1,1,1,1-2e$	
	$4,1,1,1,1,1-\{(a_1,b),(a_1,c)\}$	$2,2,1,1,1,1,1-2e$	
	$3,2,1,1,1,1-\{(a,c),e\}$		
	$3,2,1,1,1,1-\{(b,c),e\}$		
	$3,2,1,1,1,1-\{(c,d),e\}$		
	$3,2,1,1,1,1-\{(a_1,b_1),(a_1,b_2)\}$		
	$3,2,1,1,1,1-\{(a_1,b_1),(a_2,b_1)\}$		
	$2,2,2,1,1,1-2e$		
	$5,1,1,1,1,1-2e$		
	$4,2,1,1,1,1-2e$		
	$3,3,1,1,1,1-2e$		
not knotted	$2,2,1,1,1,1-\{(a,b),e\}$	$2,1,1,1,1,1,1-\{(a_1,b),(b,c)\}$	none
	$2,2,1,1,1,1-\{(c,d),e\}$	$2,1,1,1,1,1,1-\{(b,c),(d,e)\}$	
	$2,2,1,1,1,1-\{(b_1,c),(b_2,d)\}$	$2,1,1,1,1,1,1-\{(a_1,b),(a_2,c)\}$	
	$2,2,1,1,1,1-\{(a_1,c),(b_1,c)\}$	$1,1,1,1,1,1,1-2e$	
	$3,1,1,1,1,1-\{(a,b),e\}$		
	$3,1,1,1,1,1-\{(b,c),(d,e)\}$		
	$4,1,1,1,1,1-\{(a_1,b),(a_2,b)\}$		
	$4,1,1,1,1,1-\{(a_1,b),(a_2,c)\}$		
	$3,2,1,1,1,1-\{(a_1,b_1),(a_2,b_2)\}$		
	$2,1,1,1,1,1-2e$		

TABLE 9. Intrinsic knotting of 2-deficient graphs (cont.)

5. 2-DEFICIENT GRAPHS

In this section we give the classification of 2-deficient graphs with respect to intrinsic linking and knotting in Tables 5, 6, 7, 8, and 9. The argument justifying this classification is very similar to that presented in the last section and we will not include it here. See [MOR] for a full account.

Notation: We will expand the notation that we created in the last section by adding subscripts to the vertices. For example, if we are removing two edges between parts A and B of $K_{4,3,1}$, there are 3 cases to be considered: deleting two edges that share a vertex from part A ($K_{4,3,1}-\{(a_1, b_1), (a_1, b_2)\}$), deleting two edges that share a vertex from part B ($K_{4,3,1}-\{(a_1, b_1), (a_2, b_1)\}$), and deleting two edges that share no vertices ($K_{4,3,1}-\{(a_1, b_1), (a_2, b_2)\}$). Also, in some cases, if we delete a particular edge we can then delete any other without affecting the classification of the resulting graph. For example, if we delete edge (b,c) from $K_{4,3,1}$, we can then delete any other and still have an intrinsically linked graph; we will denote

graphs obtained in this way by $K_{4,3,1} - \{(b, c), e\}$. Furthermore, in some cases, we can delete any 2 edges and still have an intrinsically linked graph; for example, K_7 will be intrinsically linked no matter what 2 edges we delete. We denote these graphs $K_7\text{-}2e$.

6. GRAPHS ON 8 VERTICES

In this section we provide a classification of intrinsically knotted graphs on 8 vertices. This classification was recently independently presented by Blain, Bowlin, Fleming, Foisy, Hendricks, and LaCombe [BFHL].

Graphs on 8 vertices are subgraphs of K_8 . We will examine in turn subgraphs which are obtained by removing 1, 2, 3, ... edges from K_8 . We have already noted that K_8 , $K_8 - e$, and both $K_8 - 2e$ graphs are intrinsically knotted. Of the five graphs $K_8 - 3e$, the three which are intrinsically knotted can all be obtained by removing two edges from $K_{2,1,1,1,1,1,1}$. (See classification of 2-deficient graphs above. Note that $2, 1, 1, 1, 1, 1, 1 - \{(a, b), (c, d)\}$ and $2, 1, 1, 1, 1, 1, 1 - \{(b, c), (c, d)\}$ are the same graph.)

There are 11 graphs of the form $K_8 - 4e$. Seven of these are not knotted as they can be realized by removing an edge from one of the two unknotted $K_8 - 3e$ graphs. The remaining four are intrinsically knotted. Three of these four are of the form $K_{2,2,1,1,1,1} - 2e$. The fourth is obtained by removing 4 edges all incident to the same vertex. This graph is intrinsically knotted as it contains K_7 as a minor.

There are 24 graphs $K_8 - 5e$. All but 4 of these are not knotted as they are minors of an unknotted $K_8 - 4e$. The four intrinsically knotted $K_8 - 5e$'s are perhaps most easily described in terms of their complementary graphs. In Figure 3, Graph




FIGURE 3. Intrinsically knotted $K_8 - 5e$'s

i is $K_{3,2,1,1} - (b, c)$, or, equivalently, $K_{3,1,1,1,1,1} - \{(b, c), (c, d)\}$. The other three graphs in the figure have K_7 as a minor and are, therefore, intrinsically knotted.

Of the 56 $K_8 - 6e$ graphs, all but 6 are minors of unknotted $K_8 - 5e$'s. In Figure 4,




FIGURE 4. Intrinsically knotted $K_8 - 6e$'s

Graph i is $K_{3,2,1,1} - \{(b_1, c), (b_1, d)\}$ while Graph ii is $K_{3,3,1,1}$, or, equivalently,

$K_{3,2,1,1,1} - \{(b_1, c), (b_2, c)\}$. The remaining four graphs are obtained by splitting a vertex of K_7 and are therefore intrinsically knotted.

Only 2 of the $K_8 - 7e$ graphs are not minors of some unknotted $K_8 - 6e$. One of these two is H_8 [KS], the graph obtained by a single triangle-Y exchange on K_7 . The other is K_7 with one additional vertex. These are both intrinsically knotted. Moreover, K_7 and H_8 are minor minimal [KS]. Thus any subgraph of K_8 obtained by removing 8 or more edges is not intrinsically knotted.

In total then, there are twenty intrinsically knotted graphs on 8 vertices.

REFERENCES

- [BFHL] P. Blain, G. Bowlin, T. Fleming, J. Foisy, J. Hendricks, and J. LaCombe, ‘Some Results on Intrinsically Knotted Graphs’ (To appear in *J. Knot Theory Ramifications*).
- [B] B. Bollobás, *Extremal Graph Theory*, London Mathematical Society Monographs, **11**, Academic Press (1978).
- [CG] J.H. Conway and C.McA. Gordon, ‘Knots and Links in Spatial Graphs’, *J. Graph Theory*, **7** (1985), 445–453.
- [F1] J. Foisy, ‘Intrinsically Knotted Graphs’, *J. Graph Theory*, **39** (2002), 178–187.
- [F2] J. Foisy, ‘A Newly Recognized Intrinsically Knotted Graph’, *J. Graph Theory*, **43** (2003), 199–209.
- [F3] J. Foisy, ‘More Intrinsically Knotted Graphs’ (preprint).
- [KS] T. Kohara, S. Suzuki, ‘Some remarks on knots and links in spatial graphs’, in *Knots 90, Osaka, 1990*, de Gruyter (1992) 435–445.
- [M1] W. Mader, ‘Homomorphiesätze für Graphen’, *Math. Ann.* **178**, (1968) 154–168.
- [MOR] T.W. Mattman, R. Ottman, M. Rodrigues, ‘Intrinsic Knotting and Linking of Almost Complete Partite Graphs,’ (preprint). math.GT/0312176
- [MRS] R. Motwani, A. Raghunathan, and H. Saran, ‘Constructive Results from Graph Minors: Linkless Embeddings’, *29th Annual Symposium on Foundations of Computer Science, IEEE* (1998), 298–409.
- [NT] J. Nešetřil and R. Thomas, ‘A note on spatial representations of graphs,’ *Commentat. Math. Univ. Carolinae.* **26** (1985), 655–659.
- [OT] M. Ozawa and Y. Tsutsumi, ‘Primitive Spatial Graphs and Graph Minors,’ (preprint).
- [RS] N. Robertson and P. Seymour, ‘Graph Minors XX. Wagner’s Conjecture,’ *J. Combin. Theory Ser. B*, **92** (2004), 325–357.
- [RST] N. Robertson, P. Seymour, R. Thomas, ‘Sachs’ Linkless Embedding Conjecture’, *J. Combin. Theory Ser. B*, **64** (1995), 185–227.
- [S] H. Sachs, ‘On Spatial Representation of Finite Graphs’, in: A. Hajnal, L. Lovasz, V.T. Sós (Eds.), *Colloq. Math. Soc. János Bolyai*, North-Holland, Amsterdam, (1984), 649–662.
- [Sh] M. Shimbara, ‘Knots in certain spatial graphs,’ *Tokyo J. Math* **11** (1988), 405–413.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE UNIVERSITY, CHICO, CHICO CA 95929-0525, USA

E-mail address: Jesse.Austin.Campbell@gmail.com

E-mail address: TMattman@CSUChico.edu

DEPARTMENT OF MATHEMATICS, SOUTH HALL ROOM 6607, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, USA

E-mail address: rottman@math.ucsb.edu

MIDDLETON HIGH SCHOOL, 20932 BIG CANYON ROAD, MIDDLETOWN, CA 95461, USA

E-mail address: JDPyzer@juno.com

DEPARTMENT OF MATHEMATICS, UC BERKELEY, 970 EVANS HALL #3840, BERKELEY, CA 94720-3840, USA

E-mail address: matthewrodrigues@msn.com

DEPARTMENT OF MATHEMATICS, UNC ASHEVILLE, CPO #2350, ONE UNIVERSITY HEIGHTS,
ASHEVILLE, NC 28804-8511, USA

E-mail address: `SWillia@bulldog.unca.edu`